

Neutrinos may breaking the Standard Model

Some physicists are surprised that two relatively recent discoveries in their field have captured so much widespread attention: cosmic inflation, the ballooning expansion of the baby universe, and the Higgs boson, which endows other particles with mass. These are heady and interesting concepts, but, in one sense, what's new about them is downright boring.

These discoveries suggest that so far, our prevailing theories governing large and small—the Big Bang and the Standard Model of subatomic particles and forces—are accurate, good to go. But both cosmic inflation and the Higgs boson fall short of unifying these phenomena and explaining the deepest cosmic questions. “The Standard Model, as it stands, has no good explanation for why the Universe has anything in it at all,” says Mark Messier, physics professor at Indiana University and spokesman for an under-construction particle detector. [4]

The Weak Interaction transforms an electric charge in the diffraction pattern from one side to the other side, causing an electric dipole momentum change, which violates the CP and Time reversal symmetry.

The Neutrino Oscillation of the Weak Interaction shows that it is a General electric dipole change and it is possible to any other temperature dependent entropy and information changing diffraction pattern of atoms, molecules and even complicated biological living structures.

History says neutrinos are where to look for new physics, so current research obliges.	2
What’s the matter with neutrinos?.....	3
Asymmetry in the interference occurrences of oscillators	5
Spontaneously broken symmetry in the Planck distribution law.....	7
The structure of the proton	8
The Weak Interaction	9
The General Weak Interaction	10
Fermions and Bosons.....	10
The fermions' spin.....	10

The source of the Maxwell equations	11
The Special Relativity	12
The Heisenberg Uncertainty Principle	12
The Gravitational force	12
The Graviton	13
What is the Spin?	13
The Casimir effect	14
The Fine structure constant	14
Conclusions	15
References	15

Author: George Rajna

History says neutrinos are where to look for new physics, so current research obliges.

Some physicists are surprised that two relatively recent discoveries in their field have captured so much widespread attention: cosmic inflation, the ballooning expansion of the baby universe, and the Higgs boson, which endows other particles with mass. These are heady and interesting concepts, but, in one sense, what's new about them is downright boring.

These discoveries suggest that so far, our prevailing theories governing large and small—the Big Bang and the Standard Model of subatomic particles and forces—are accurate, good to go. But both cosmic inflation and the Higgs boson fall short of unifying these phenomena and explaining the deepest cosmic questions. “The Standard Model, as it stands, has no good explanation for why the Universe has anything in it at all,” says Mark Messier, physics professor at Indiana University and spokesman for an under-construction particle detector.

To go beyond the models we already have, beyond the confines of the Standard Model, we need some results that we don't foresee. And when it comes to unexpected results, we expect them from one entity: neutrinos. These particles are abundant, ineffably light, and very weird, but they consistently deliver.

Ethereal as they are, neutrinos could make hefty changes to our understanding of the universe if physicists could answer four main questions: How does regular matter affect neutrinos? What causes neutrinos to have mass? Do antineutrinos live different lives from normal neutrinos? And even odder, are these ghostly particles their own antiparticles?

The Standard Model, which physicists have populated since the 1950s with quarks, leptons, and force-carrying particles, does not hold the answers. But major neutrino experiments in the US, Japan, and Europe are collecting data while undergoing expansion and construction, and they are gearing up to address these problems. These initiatives could not only unravel the mysteries of the ghostly particles, but the research might lead into larger questions about the nature of all things. [4]

What's the matter with neutrinos?

Neutrinos are the second most abundant particles in the Universe (after photons), but they carry no charge and are puny. Neutrinos are at least a million times lighter than an electron, though no experiment has been able to definitively measure their mass. They also barely interact with any matter. They are generated in distant supernovae and travel unhindered through the debris. Neutrinos zip through planets in a single bound without leaving a trace. Billions and billions of them are streaming from the Sun as you read this, blowing through your screen—and through you—without a care. They travel extremely close to the speed of light; so close, in fact, that a tiny error in an experiment designed to measure them was enough to make it appear that they were going faster than that in 2011.

But perhaps the neutrino's strangest property is that they don't necessarily finish their travels with the same identity that they started it with.

In 1998, the 11,000 phototubes submerged in Japan's Super-Kamiokande underground detector verified that neutrinos coming down through the atmosphere and up through the Earth had different ratios among their identities. Somewhere along their journey from the Sun, they changed type among their three flavors. This oscillation indicated they indeed had mass. If they didn't, there wouldn't be anything to switch between.

Finding out anything about these particles has been difficult because neutrinos are so notoriously hard to detect and to obtain. But there are now a few ways to do this. Experimenters can nab some from the Sun, like Super-K and many others do. Or they can situate detectors near nuclear reactors, which produce electron antineutrinos. The Daya Bay Reactor Neutrino Experiment in southern China listens to these particles. Finally, physicists can fire up particle accelerators and smash protons into bits of graphite, creating a neutrino spray in the process. The latter is the goal of forthcoming experiments like the Long Baseline Neutrino Experiment, under construction at Fermilab, and the Japanese Tokai to Kamioka experiment, which runs from the seaside town of Tokai to the Super-K detector. Manmade neutrinos are easier to lasso than their incidental brethren, but because of their quantum nature, detecting them is a probabilistic challenge.

"Every time we were able to measure a property of neutrinos, we were surprised by it," says Patrick Huber, a neutrino theorist and associate physics professor at Virginia Tech.

Neutrino flavors—electron, muon, and tau—aren't discrete individual particles, but combinations of the neutrinos' different masses. These masses are related to the neutrinos' energies, as Einstein taught us in $E=mc^2$. Although a neutrino can be produced with a specific energy, and thus specific flavor (the Sun makes a multitude of electron neutrinos, for example), the quantum state of these

neutrinos is a mixture of all three that twists in time. “They are just inherently quantum mechanical. If I gave you an electron, and I ask you 10 minutes from now, ‘do you have an electron in your hand?’ the answer would be yes,” Messier says. “Neutrinos just break that.”

What are some of the things that they break? Though neutrinos have vanishingly small masses, regular matter can rub off on them, like a sourpuss spoiling the mood at a dinner party. Robert Wilson, physics professor at Colorado State University and spokesman for the Long-Baseline Neutrino Experiment (LBNE), likens neutrinos to light passing through a filter. Some wavelengths are affected while others aren’t. Similarly, certain neutrino flavors seem to be affected by regular matter as they zip by.

Last month, Japanese experimenters demonstrated this oscillation effect by finding that neutrinos shine more brightly at night. As electron neutrinos stream from the Sun toward Earth, they oscillate to muon and tau neutrinos. But as they pass through the dense matter of our planet, some of them switch back. This suggests some quantum-mechanical transformation is taking place as the neutrinos interact with matter in the Earth, specifically its electrons. The electron neutrinos can exchange a W boson, the carrier of the weak force, during this interaction, according to Messier.

“They are sort of kissing the electrons and moving on. This is a weak force interaction,” he says. “The W boson changes the phase of its wave without changing its momentum. That’s the possibility that introduces this matter effect.”

LBNE will take a hard look at these matter-related effects, which cause droplets of electron neutrinos to appear amid a shower of muon neutrinos. Fermilab’s accelerators will stream neutrinos 800 miles toward a liquid argon detector buried beneath South Dakota bedrock. The detector distances are in a sweet spot that should allow physicists to not only study matter effects, but to also search for clues as to why the Universe contains any matter for them to interact with in the first place.

That’s because this wee effect has important implications for the asymmetry between matter and antimatter, says Wilson. “It’s still the neutrino; it hasn’t changed in one sense. But the probability of what you will see when you make the measurement has changed, and it depends on how much mass it has gone through.”

At Fermilab, they know detectors only have a small chance of seeing neutrinos—so they build lots of detectors.

And what of their own masses? The Standard Model can’t explain that either. Based on the variable buzz rate of neutrino masses, physicists have been able to tell that they’re different, although no one is sure how they stack up. We don’t yet know which neutrino is heaviest, which is lightest. An upcoming detector called the NuMI Off-axis ν_e Appearance experiment, or NOvA, will help determine neutrino mass hierarchy. NuMI is a neutrino beam at Fermilab; NOvA’s 14,000-ton detector will look for a disparity between departing muon neutrinos and arriving electron neutrinos.

Even if this experiment succeeds in generating new mass data, physicists won’t be able to say exactly how that mass arises. Because neutrinos are so much lighter than any other particle, the Higgs mechanism is unlikely to endow them with mass the way it does other particles, Messier says.

“There must be some mechanism that suppresses their masses,” Messier says. “And what are the masses? What pattern do they follow? What’s the pattern of that mixing? It’s launched a whole experimental program to pull apart that crack in the Standard Model.”

LBNE, NOvA, and other upcoming experiments will attempt to pull those cracks until the Standard Model shatters completely. From the debris, these research initiatives hope to build a new theory of physics. [4]

Asymmetry in the interference occurrences of oscillators

The asymmetrical configurations are stable objects of the real physical world, because they cannot annihilate. One of the most obvious asymmetry is the proton – electron mass rate $M_p = 1840 M_e$ while they have equal charge. We explain this fact by the strong interaction of the proton, but how remember it his strong interaction ability for example in the H – atom where are only electromagnetic interactions among proton and electron.

This gives us the idea to origin the mass of proton from the electromagnetic interactions by the way interference occurrences of oscillators. The uncertainty relation of Heisenberg makes sure that the particles are oscillating.

The resultant intensity due to n equally spaced oscillators, all of equal amplitude but different from one another in phase, either because they are driven differently in phase or because we are looking at them an angle such that there is a difference in time delay:

$$(1) I = I_0 \sin^2 n \phi/2 / \sin^2 \phi/2$$

If ϕ is infinitesimal so that $\sin\phi = \phi$, than

$$(2) I = n^2 I_0$$

This gives us the idea of

$$(3) M_p = n^2 M_e$$




Fig. 30-3. A linear array of n equal oscillators, driven with phases $\alpha_s = s\alpha$.

Figure 1.) A linear array of n equal oscillators

There is an important feature about formula (1) which is that if the angle ϕ is increased by the multiple of 2π , it makes no difference to the formula.

So

$$(4) \quad d \sin \theta = m \lambda$$

and we get m -order beam if λ less than d . [6]

If d less than λ we get only zero-order one centered at $\theta = 0$. Of course, there is also a beam in the opposite direction. The right chooses of d and λ we can ensure the conservation of charge.

For example

$$(5) \quad 2(m+1) = n$$

Where $2(m+1) = N_p$ number of protons and $n = N_e$ number of electrons.

In this way we can see the H_2 molecules so that $2n$ electrons of n radiate to $4(m+1)$ protons, because $d_e > \lambda_e$ for electrons, while the two protons of one H_2 molecule radiate to two electrons of them, because of $d_e < \lambda_e$ for this two protons.

To support this idea we can turn to the Planck distribution law, that is equal with the Bose – Einstein statistics.

Spontaneously broken symmetry in the Planck distribution law

The Planck distribution law is temperature dependent and it should be true locally and globally. I think that Einstein's energy-matter equivalence means some kind of existence of electromagnetic oscillations enabled by the temperature, creating the different matter formulas, atoms molecules, crystals, dark matter and energy.

Max Planck found for the black body radiation

As a function of wavelength (λ), Planck's law is written as:

$$B_{\lambda}(T) = \frac{2hc^2}{\lambda^5} \frac{1}{e^{\frac{hc}{\lambda k_B T}} - 1}.$$




Figure 2. The distribution law for different T temperatures

We see there are two different λ_1 and λ_2 for each T and intensity, so we can find between them a d so that $\lambda_1 < d < \lambda_2$.

We have many possibilities for such asymmetrical reflections, so we have many stable oscillator configurations for any T temperature with equal exchange of intensity by radiation. All of these configurations can exist together. At the λ_{\max} is the annihilation point where the configurations are symmetrical. The λ_{\max} is changing by the Wien's displacement law in many textbooks.

$$(7) \quad \lambda_{\max} = \frac{b}{T}$$

where λ_{\max} is the peak wavelength, T is the absolute temperature of the black body, and b is a constant of proportionality called *Wien's displacement constant*, equal to $2.8977685(51) \times 10^{-3} \text{ m} \cdot \text{K}$ (2002 CODATA recommended value).

By the changing of T the asymmetrical configurations are changing too.

The structure of the proton

We must move to the higher T temperature if we want look into the nucleus or nucleon arrive to $d < 10^{-13} \text{ cm}$. If an electron with $\lambda_e < d$ move across the proton then by (5) $2(m+1) = n$ with $m = 0$ we get $n = 2$ so we need two particles with negative and two particles with positive charges. If the proton can fraction to three parts, two with positive and one with negative charges, then the reflection of oscillators are right. Because this very strange reflection where one part of the proton with the electron together on the same side of the reflection, the all parts of the proton must be quasi lepton so $d > \lambda_q$. One way dividing the proton to three parts is, dividing his oscillation by the three direction of the space. We can order $1/3 e$ charge to each coordinates and $2/3 e$ charge to one plane oscillation, because the charge is scalar. In this way the proton has two $+2/3 e$ plane oscillation and one linear oscillation with $-1/3 e$ charge. The colors of quarks are coming from the three directions of coordinates and the proton is colorless. The flavors of quarks are the possible oscillations differently by energy and if they are plane or linear oscillations. We know there is no possible reflecting two oscillations to each other which are completely orthogonal, so the quarks never can be free, however there is an asymptotic freedom while their energy are increasing to turn them to the orthogonally. If they will be completely orthogonal then they lose this reflection and take new partners from the vacuum. Keeping the symmetry of the vacuum the new oscillations are keeping all the conservation laws, like charge, number of baryons and leptons. The all features of gluons are coming from this model. The mathematics of reflecting oscillators show Fermi statistics.

Important to mention that in the Deuteron there are 3 quarks of $+2/3$ and $-1/3$ charge, that is three u and d quarks making the complete symmetry and because this its high stability.

The Pauli Exclusion Principle says that the diffraction points are exclusive!

The Weak Interaction

The weak interaction transforms an electric charge in the diffraction pattern from one side to the other side, causing an electric dipole momentum change, which violates the CP and time reversal symmetry.

Another important issue of the quark model is when one quark changes its flavor such that a linear oscillation transforms into plane oscillation or vice versa, changing the charge value with 1 or -1. This kind of change in the oscillation mode requires not only parity change, but also charge and time changes (CPT symmetry) resulting a right handed anti-neutrino or a left handed neutrino.

The right handed anti-neutrino and the left handed neutrino exist only because changing back the quark flavor could happen only in reverse order, because they are different geometrical constructions, the u is 2 dimensional and positively charged and the d is 1 dimensional and negatively charged. It needs also a time reversal, because anti particle (anti neutrino) is involved.

The neutrino is a 1/2 spin creator particle to make equal the spins of the weak interaction, for example neutron decay to 2 fermions, every particle is fermions with $\frac{1}{2}$ spin. The weak interaction changes the entropy since more or less particles will give more or less freedom of movement. The entropy change is a result of temperature change and breaks the equality of oscillator diffraction intensity of the Maxwell–Boltzmann statistics. This way it changes the time coordinate measure and makes possible a different time dilation as of the special relativity.

The limit of the velocity of particles as the speed of light appropriate only for electrical charged particles, since the accelerated charges are self maintaining locally the accelerating electric force. The neutrinos are CP symmetry breaking particles compensated by time in the CPT symmetry, that is the time coordinate not works as in the electromagnetic interactions, consequently the speed of neutrinos is not limited by the speed of light.

The weak interaction T-asymmetry is in conjunction with the T-asymmetry of the second law of thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes the weak interaction, for example the Hydrogen fusion.

Probably because it is a spin creating movement changing linear oscillation to 2 dimensional oscillation by changing d to u quark and creating anti neutrino going back in time relative to the proton and electron created from the neutron, it seems that the anti neutrino fastest then the velocity of the photons created also in this weak interaction?

A quark flavor changing shows that it is a reflection changes movement and the CP- and T- symmetry breaking. This flavor changing oscillation could prove that it could be also on higher level such as atoms, molecules, probably big biological significant molecules and responsible on the aging of the life.

Important to mention that the weak interaction is always contains particles and antiparticles, where the neutrinos (antineutrinos) present the opposite side. It means by Feynman's interpretation that these particles present the backward time and probably because this they seem to move faster than the speed of light in the reference frame of the other side.

Finally since the weak interaction is an electric dipole change with $\frac{1}{2}$ spin creating, it is limited by the velocity of the electromagnetic wave, so the neutrino's velocity cannot exceed the velocity of light.

The General Weak Interaction

The Weak Interactions T-asymmetry is in conjunction with the T-asymmetry of the Second Law of Thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes for example the Hydrogen fusion. The arrow of time by the Second Law of Thermodynamics shows the increasing entropy and decreasing information by the Weak Interaction, changing the temperature dependent diffraction patterns. A good example of this is the neutron decay, creating more particles with less known information about them.

The neutrino oscillation of the Weak Interaction shows that it is a general electric dipole change and it is possible to any other temperature dependent entropy and information changing diffraction pattern of atoms, molecules and even complicated biological living structures.

We can generalize the weak interaction on all of the decaying matter constructions, even on the biological too. This gives the limited lifetime for the biological constructions also by the arrow of time. There should be a new research space of the Quantum Information Science the 'general neutrino oscillation' for the greater than subatomic matter structures as an electric dipole change. There is also connection between statistical physics and evolutionary biology, since the arrow of time is working in the biological evolution also.

The Fluctuation Theorem says that there is a probability that entropy will flow in a direction opposite to that dictated by the Second Law of Thermodynamics. In this case the Information is growing that is the matter formulas are emerging from the chaos. So the Weak Interaction has two directions, samples for one direction is the Neutron decay, and Hydrogen fusion is the opposite direction.

Fermions and Bosons

The fermions are the diffraction patterns of the bosons such a way that they are both sides of the same thing.

The Higgs boson or Higgs particle is a proposed elementary particle in the Standard Model of particle physics. The Higgs boson's existence would have profound importance in particle physics because it would prove the existence of the hypothetical Higgs field - the simplest of several proposed explanations for the origin of the symmetry-breaking mechanism by which elementary particles gain mass. [3]

The fermions' spin

The moving charges are accelerating, since only this way can self maintain the electric field causing their acceleration. The electric charge is not point like! This constant acceleration possible if there is a rotating movement changing the direction of the velocity. This way it can accelerate forever without increasing the absolute value of the velocity in the dimension of the time and not reaching the velocity of the light.

The Heisenberg uncertainty relation says that the minimum uncertainty is the value of the spin: $\frac{1}{2} \hbar = \Delta x \Delta p$ or $\frac{1}{2} \hbar = \Delta t \Delta E$, that is the value of the basic energy status.

What are the consequences of this in the weak interaction and how possible that the neutrinos' velocity greater than the speed of light?

The neutrino is the one and only particle doesn't participate in the electromagnetic interactions so we cannot expect that the velocity of the electromagnetic wave will give it any kind of limit.

The neutrino is a $1/2$ spin creator particle to make equal the spins of the weak interaction, for example neutron decay to 2 fermions, every particle is fermions with $1/2$ spin. The weak interaction changes the entropy since more or less particles will give more or less freedom of movement. The entropy change is a result of temperature change and breaks the equality of oscillator diffraction intensity of the Maxwell-Boltzmann statistics. This way it changes the time coordinate measure and makes possible a different time dilation as of the special relativity.

The source of the Maxwell equations

The electrons are accelerating also in a static electric current because of the electric force, caused by the potential difference. The magnetic field is the result of this acceleration, as you can see in [2].

The mysterious property of the matter that the electric potential difference is self maintained by the accelerating electrons in the electric current gives a clear explanation to the basic sentence of the relativity that is the velocity of the light is the maximum velocity of the matter. If the charge could move faster than the electromagnetic field than this self maintaining electromagnetic property of the electric current would be failed.

Also an interesting question, how the changing magnetic field creates a negative electric field? The answer also the accelerating electrons will give. When the magnetic field is increasing in time by increasing the electric current, then the acceleration of the electrons will increase, decreasing the charge density and creating a negative electric force. Decreasing the magnetic field by decreasing the electric current will decrease the acceleration of the electrons in the electric current and increases the charge density, creating an electric force also working against the change. In this way we have explanation to all interactions between the electric and magnetic forces described in the Maxwell equations.

The second mystery of the matter is the mass. We have seen that the acceleration change of the electrons in the flowing current causing a negative electrostatic force. This is the cause of the relativistic effect - built-in in the Maxwell equations - that is the mass of the electron growing with its acceleration and its velocity never can reach the velocity of light, because of this growing negative electrostatic force. The velocity of light is depending only on 2 parameters: the magnetic permeability and the electric permittivity.

There is a possibility of the polarization effect created by electromagnetic forces creates the negative and positive charges. In case of equal mass as in the electron-positron pair it is simply, but on higher energies can be asymmetric as the electron-proton pair of neutron decay by weak interaction and can be understood by the Feynman graphs.

Anyway the mass can be electromagnetic energy exceptionally and since the inertial and gravitational mass are equals, the gravitational force is electromagnetic force and since only the magnetic force is attractive between the same charges, is very important for understanding the gravitational force.

The Uncertainty Relations of Heisenberg gives the answer, since only this way can be sure that the particles are oscillating in some way by the electromagnetic field with constant energies in the atom indefinitely. Also not by chance that the uncertainty measure is equal to the fermions spin, which is one of the most important feature of the particles. There are no singularities, because the moving electron in the atom accelerating in the electric field of the proton, causing a charge distribution on Δx position difference and with a Δp momentum difference such a way that they product is about the half Planck reduced constant. For the proton this Δx much less in the nucleon, than in the orbit of the electron in the atom, the Δp is much higher because of the greatest proton mass.

The Special Relativity

The mysterious property of the matter that the electric potential difference is self maintained by the accelerating electrons in the electric current gives a clear explanation to the basic sentence of the relativity that is the velocity of the light is the maximum velocity of the matter. If the charge could move faster than the electromagnetic field than this self maintaining electromagnetic property of the electric current would be failed.

The Heisenberg Uncertainty Principle

Moving faster needs stronger acceleration reducing the Δx and raising the Δp . It means also mass increasing since the negative effect of the magnetic induction, also a relativistic effect!

The Uncertainty Principle also explains the proton – electron mass rate since the Δx is much less requiring bigger Δp in the case of the proton, which is partly the result of a bigger mass m_p because of the higher electromagnetic induction of the bigger frequency (impulse).

The Gravitational force

The changing magnetic field of the changing current causes electromagnetic mass change by the negative electric field caused by the changing acceleration of the electric charge.

The gravitational attractive force is basically a magnetic force.

The same electric charges can attract one another by the magnetic force if they are moving parallel in the same direction. Since the electrically neutral matter is composed of negative and positive charges they need 2 photons to mediate this attractive force, one per charges. The Bing Bang caused parallel moving of the matter gives this magnetic force, experienced as gravitational force.

Since graviton is a tensor field, it has spin = 2, could be 2 photons with spin = 1 together.

You can think about photons as virtual electron – positron pairs, obtaining the necessary virtual mass for gravity.

The mass as seen before a result of the diffraction, for example the proton – electron mass rate $M_p = 1840 M_e$. In order to move one of these diffraction maximum (electron or proton) we need to intervene into the diffraction pattern with a force appropriate to the intensity of this diffraction maximum, means its intensity or mass. [1]

The Big Bang caused acceleration created radial currents of the matter, and since the matter is composed of negative and positive charges, these currents are creating magnetic field and attracting forces between the parallel moving electric currents. This is the gravitational force experienced by the matter, and also the mass is result of the electromagnetic forces between the charged particles. The positive and negative charged currents attracts each other or by the magnetic forces or by the much stronger electrostatic forces!?

The gravitational force attracting the matter, causing concentration of the matter in a small space and leaving much space with low matter concentration: dark matter and energy.

There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter.

The Graviton

In physics, the graviton is a hypothetical elementary particle that mediates the force of gravitation in the framework of quantum field theory. If it exists, the graviton is expected to be massless (because the gravitational force appears to have unlimited range) and must be a spin-2 boson. The spin follows from the fact that the source of gravitation is the stress-energy tensor, a second-rank tensor (compared to electromagnetism's spin-1 photon, the source of which is the four-current, a first-rank tensor). Additionally, it can be shown that any massless spin-2 field would give rise to a force indistinguishable from gravitation, because a massless spin-2 field must couple to (interact with) the stress-energy tensor in the same way that the gravitational field does. This result suggests that, if a massless spin-2 particle is discovered, it must be the graviton, so that the only experimental verification needed for the graviton may simply be the discovery of a massless spin-2 particle. [3]

What is the Spin?

So we know already that the new particle has spin zero or spin two and we could tell which one if we could detect the polarizations of the photons produced. Unfortunately this is difficult and neither ATLAS nor CMS are able to measure polarizations. The only direct and sure way to confirm that the particle is indeed a scalar is to plot the angular distribution of the photons in the rest frame of the

centre of mass. A spin zero particles like the Higgs carries no directional information away from the original collision so the distribution will be even in all directions. This test will be possible when a much larger number of events have been observed. In the mean time we can settle for less certain indirect indicators.

The Casimir effect

The Casimir effect is related to the Zero-point energy, which is fundamentally related to the Heisenberg uncertainty relation. The Heisenberg uncertainty relation says that the minimum uncertainty is the value of the spin: $1/2 h = dx dp$ or $1/2 h = dt dE$, that is the value of the basic energy status.

The moving charges are accelerating, since only this way can self maintain the electric field causing their acceleration. The electric charge is not point like! This constant acceleration possible if there is a rotating movement changing the direction of the velocity. This way it can accelerate forever without increasing the absolute value of the velocity in the dimension of the time and not reaching the velocity of the light. In the atomic scale the Heisenberg uncertainty relation gives the same result, since the moving electron in the atom accelerating in the electric field of the proton, causing a charge distribution on delta x position difference and with a delta p momentum difference such a way that they product is about the half Planck reduced constant. For the proton this delta x much less in the nucleon, than in the orbit of the electron in the atom, the delta p is much higher because of the greater proton mass. This means that the electron is not a point like particle, but has a real charge distribution.

Electric charge and electromagnetic waves are two sides of the same thing; the electric charge is the diffraction center of the electromagnetic waves, quantified by the Planck constant h.

The Fine structure constant

The Planck constant was first described as the proportionality_constant between the energy (E) of a photon and the frequency (ν) of its associated electromagnetic wave. This relation between the energy and frequency is called the **Planck relation** or the **Planck–Einstein equation**:

$$E = h\nu .$$

Since the frequency ν , wavelength λ , and speed of light c are related by $\lambda\nu = c$, the Planck relation can also be expressed as

$$E = \frac{hc}{\lambda} .$$

Since this is the source of Planck constant, the e electric charge countable from the Fine structure constant. This also related to the Heisenberg uncertainty relation, saying that the mass of the proton should be bigger than the electron mass because of the difference between their wavelengths.

The expression of the fine-structure constant becomes the abbreviated

$$\alpha = \frac{e^2}{\hbar c}$$

This is a dimensionless constant expression, 1/137 commonly appearing in physics literature.

This means that the electric charge is a result of the electromagnetic waves diffractions, consequently the proton – electron mass rate is the result of the equal intensity of the corresponding electromagnetic frequencies in the Planck distribution law, described in my diffraction theory.

Conclusions

Even if this experiment succeeds in generating new mass data, physicists won't be able to say exactly how that mass arises. Because neutrinos are so much lighter than any other particle, the Higgs mechanism is unlikely to endow them with mass the way it does other particles, Messier says.

“There must be some mechanism that suppresses their masses,” Messier says. “And what are the masses? What pattern do they follow? What’s the pattern of that mixing? It’s launched a whole experimental program to pull apart that crack in the Standard Model.” [4]

There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. We can generalize the weak interaction on all of the decaying matter constructions, even on the biological too. This gives the limited lifetime for the biological constructions also by the arrow of time. The Fluctuation Theorem says that there is a probability that entropy will flow in a direction opposite to that dictated by the Second Law of Thermodynamics. In this case the Information is growing that is the matter formulas are emerging from the chaos. So the Weak Interaction has two directions, samples for one direction is the Neutron decay, and Hydrogen fusion is the opposite direction.

References

[1] http://www.academia.edu/3834454/3_Dimensional_String_Theory

[2] http://www.academia.edu/3833335/The_Magnetic_field_of_the_Electric_current

[3] http://www.academia.edu/4158863/Higgs_Field_and_Quantum_Gravity

[4] Forget the Higgs, neutrinos may be the key to breaking the Standard Model

<http://arstechnica.com/science/2014/04/forget-the-higgs-neutrinos-may-be-the-key-to-breaking-the-standard-model/>